

Absolute continuity of autophage measures on finite-dimensional vector spaces

C. R. E. Raja

Abstract

We consider a class of measures called autophage which was introduced and studied by Szekely for measures on the real line. We show that the autophage measures on finite-dimensional vector spaces over real or $\mathbb{Q}_p$ are infinitely divisible without idempotent factors and are absolutely continuous with bounded continuous density. We also show that certain semistable measures on such vector spaces are absolutely continuous.

1 Introduction and Preliminaries

Let G be a locally compact second countable group. Let $\mathcal{P}(G)$ be the space of Borel probability measures on G , equipped with the weak topology. For any μ and $\lambda \in \mathcal{P}(G)$, $\mu * \lambda$ denotes the convolution of λ and μ . For $\mu \in \mathcal{P}(G)$, $\check{\mu}$ is defined by $\check{\mu}(B) = \mu(B^{-1})$ for any Borel subset B of G and for any integer $n \geq 1$, μ^n denotes the n th-convolution of power of μ . We say that a measure $\mu \in \mathcal{P}(G)$ is *continuous* if $\mu(\{x\}) = 0$ for all $x \in G$ and *absolutely continuous* if μ is absolutely continuous with respect to a Haar measure on G . A *continuous convolution semigroup* (μ_t) in $\mathcal{P}(G)$ is a continuous homomorphism $t \mapsto \mu_t$ from the additive semigroup $\mathbb{R}^+ \cup 0$ into $\mathcal{P}(G)$.

Let $\text{Aut}(G)$ denotes the group of all continuous automorphisms of G . Then for any $T \in \text{Aut}(G)$ and $\lambda \in \mathcal{P}(G)$, $T(\lambda)$ is defined by $T(\lambda)(B) = \lambda(T^{-1}(B))$ for any Borel subset B of G . An automorphism $T \in \text{Aut}(G)$ is called *contraction* if $T^n x \rightarrow e$ for all $x \in G$. In case, G is a finite-dimensional vector space over real or $\mathbb{Q}_p$, we say that $T \in \text{Aut}(G)$ is a *strict-contraction* if $\|T\| < 1$, where $\|\cdot\|$ is the operator norm.

We say that a $\lambda \in \mathcal{P}(G)$ is a factor of a $\mu \in \mathcal{P}(G)$ if there exists a $\nu \in \mathcal{P}(G)$ such that $\mu = T(\lambda) * \nu$ and ν is called co-factor. Here we wish to study the measures μ for which there exists commuting contractions T and S such that $T(\mu)$ is a factor of μ with $S(\mu)$ as the corresponding co-factor. This type of measures on the real line was introduced and studied by Szekely and Szekely calls such measures autophage

(see [10]). He also proved that nondegenerate autophage measures are infinitely divisible and absolutely continuous with a bounded continuous density. The main part of his proof was to prove that the Fourier transform of such measures are integrable. This implies by Fourier inversion formula that such measures are absolutely continuous with continuous bounded density. Here we prove that autophage measures on finite dimensional vector spaces over real or $\mathbb{Q}_p$ are infinitely divisible and absolutely continuous. For a given infinitely divisible probability measure it is of considerable interest to know when it is absolutely continuous. In this point of view these results are of considerable interest. We also show that certain semistable measures on vector spaces over real and $\mathbb{Q}_p$ are absolutely continuous with bounded continuous density: see [3] for results on the absolute continuity of semistable measures on $\mathbb{R}^N$.

2 Menu-card of measures

Definition 2.1 We say that a measure $\nu \in \mathcal{P}(G)$ belongs to the *menu-card* of a measure $\mu \in \mathcal{P}(G)$ if $\mu = T(\mu) * \nu$ for some contraction T of G .

Remark 2.1 It is easy to see that a menu-card of a measure μ is non-empty if and only if μ is strongly T -decomposable for some contraction T : a measure $\mu \in \mathcal{P}(G)$ is strongly T -decomposable for a $T \in \text{Aut}(G)$ if $T(\mu)$ is a factor of μ and $T^n(\mu) \rightarrow \delta_e$ (see [9] and [6] for results regarding strongly T -decomposable measures).

We now prove that the nondegenerate strongly T -decomposable measures on real and p -adic unipotent groups, studied in [6] are continuous which implies the continuity of measures with nontrivial menu-card. By an algebraic group over real or $\mathbb{Q}_p$ we mean the group of real or $\mathbb{Q}_p$ points of an algebraic group defined over real or $\mathbb{Q}_p$ respectively.

Proposition 2.1 *Let G be a unipotent algebraic group over real or $\mathbb{Q}_p$. Then nondegenerate strongly T -decomposable measures on G are diffuse, that is, continuous. In particular, if the menu-card of a measure $\mu \in \mathcal{P}(G)$ contains a nondegenerate measure then μ is continuous.*

Proof Let $T \in \text{Aut}(G)$ be such that $\mu = T(\mu) * \nu$ for some $\nu \in \mathcal{P}(G)$ and $T^n(\mu) \rightarrow \delta_e$. Suppose μ is not continuous. We now claim that μ degenerates. Let Z be the center of G . Let $G_0 = G$ and for $i \geq 1$, $G_i = [G, G_{i-1}]$. There exists a $k \geq 0$ such that $G_k \neq (e)$ but $G_{k+1} = (e)$. Proof is based on induction on k . Suppose $k = 0$, then G is abelian. Since G has no elements of finite order, by 3.5 of [9], ν degenerates and hence μ degenerates. Suppose $k > 0$. Let $G' = G/Z$ and μ' and ν' be the projection of μ and ν under the canonical map from G to G' . Let T' be the factor automorphism

of G' corresponding to T . We then have $\mu' = T'\mu' * \nu'$. It may be easily seen that μ' is also not continuous and hence by induction μ' as well as ν' degenerates. Now, let $\lambda = \mu * \check{\mu}$ and $\rho = \nu * \check{\nu}$. Then $\lambda = T(\mu) * \nu * \check{\nu} * T(\check{\mu}) = T(\lambda) * \rho$ since $\nu * \check{\nu}$ is supported on the center of G . Also, λ and ρ are supported on the center of G and the center of G is T -invariant. Since μ is not continuous, there exists a $x_0 \in G$ such that $\mu(\{x_0\}) > 0$. Now, $\lambda(\{e\}) = \mu * \check{\mu}(\{e\}) \geq \mu(\{x_0\})^2 > 0$. Thus, λ is also not continuous. Thus, we infer from the abelian case that λ degenerates and so does μ .

3 Autophage measure

Definition 3.1 A measure $\mu \in \mathcal{P}(G)$ is called *autophage* if there exists contractions T and S of G such that $\mu = T(\mu) * S(\mu)$ and $TS = ST$.

In [10] and [7] autophage measures were introduced on the real line $\mathbb{R}$. We recall the definition of autophage on $\mathbb{R}$, $\mu \in \mathcal{P}(\mathbb{R})$ is autophage if $F(x) = F(ax)F(bx)$ for some $a, b > 1$ where F is the distribution of μ . Our generalization in Definition 3.1 for measures on general groups, we replace the homotheties by two commuting contractions. If we had chosen two elements of a one-parameter semigroup of contractions in place of homotheties, our notion would be very restrictive. Because in the p -adic situation, there are no one-parameter groups but there are many autophage measures in the sense of Definition 3.1. Thus, our choice of commuting contractions in Definition 3.1 yields more autophage measures.

It may be easily seen that stable measures on finite-dimensional vector spaces are autophage. More generally, measures that are stable with respect to a commutative group of contractive automorphisms are autophage. We recall that a measure μ in $\mathcal{P}(G)$ is said to be (T, n) -semistable for some $n \in \mathbb{N}$ and for an automorphism T if $T(\mu^n) = \mu$. It may be easily seen that $(T, 2)$ -semistable measure is autophage when T is a contractive automorphism.

We now directly show that Gaussian measures on $\mathbb{R}^n$ are autophage. Let μ be a symmetric Gaussian measure on $\mathbb{R}^n$. Then $\hat{\mu}(x) = \exp - \langle Px, x \rangle$ where P is a positive definite matrix. Let T be any strict-contraction on $\mathbb{R}^n$ commuting with P such that T is invertible and symmetric (we may choose T to be a non-zero homothesy). Then for any $x \in \mathbb{R}^n$, $0 < \|x\|^2 - \|T(x)\|^2 = \langle I - T^2(x), x \rangle$, that is, $I - T^2$ is positive definite. Let S be the unique square root of $I - T^2$. Then S is invertible and commutes with T . Since T commutes with P , S also commutes with P . This implies that $TPT^* + SPS^* = T^2P + S^2P = (T^2 + S^2)P = P$. Thus, $\mu = T(\mu) * S(\mu)$ where T and S are mutually commuting strict-contractions.

4 Infinite divisibility

For any two contractions T and S on G , let $sg(T, S)$ be the semigroup generated by T and S . For any $\alpha \in sg(T, S)$, define $l(\alpha)$ to be the sum of number of T and S appear in the expression of α . Then we have

Lemma 4.1 *Let V be a finite-dimensional vector space over real or $\mathbb{Q}_p$. Let T and S be strict-contractions on V . Then $\alpha(x) \rightarrow e$ for all $x \in V$ as $l(\alpha) \rightarrow \infty$.*

Proof Since T and S are strict-contractions, for a given $\epsilon > 0$, there exists a $k \geq 1$, $\|T\|^n < 1$ and $\|S\|^n < 1$ for all $n \geq k$. For $\alpha \in sg(T, S)$ with $l(\alpha) = n$, we have $\|\alpha\| \leq \|T\|^{n-i} \|S\|^i$ for some $0 \leq i \leq n$. If $n = l(\alpha) \geq 2k$, then $\|\alpha\| < \epsilon$. Thus, $\|\alpha\| \rightarrow 0$ as $l(\alpha) \rightarrow \infty$.

Proposition 4.1 *Let V be a finite-dimensional vector space over real or $\mathbb{Q}_p$. Let $T_1, T_2, \dots, T_m$ be strict-contractions on V . Suppose μ is a probability measure such that $\mu = T_1(\mu) * T_2(\mu) * \dots * T_m(\mu)$. Then μ is infinitely divisible without idempotent factors.*

Proof We treat the case $m = 2$ and the general case may be done in a similar fashion. Let T and S be strict-contractions on G such that $\mu = T(\mu) * S(\mu)$. We first claim that μ is infinitely divisible. Iteratively we get that

$$\begin{aligned} \mu &= T(\mu)S(\mu) \\ &= T^2(\mu) * TS(\mu) * ST(\mu) * S^2(\mu) \\ &= T^3(\mu) * T^2S(\mu) * TST(\mu) * TS^2(\mu) * ST^2(\mu) * STS(\mu) * S^2T(\mu) * S^3(\mu) \\ &= \dots\dots\dots \\ &= \prod_{\alpha} \alpha(\mu) \end{aligned}$$

where the product varies over all α in $\{\alpha \in sg(T, S) \mid l(\alpha) = n\}$. Thus, we have a triangular system of probability measure where sum of the row product is μ and the n th row is

$$T^n(\mu), T^{n-1}S(\mu), \dots, S^{n-1}T(\mu), S^n(\mu).$$

Since the group is abelian, it is enough to claim that the triangular system is infinitesimal. That is, we should prove that $\alpha(\mu) \rightarrow e$ as $l(\alpha) \rightarrow \infty$ which follows from Lemma 4.1 and dominated convergence theorem.

We now claim that μ has no idempotent factors. We may assume that V is a p -adic vector space. Let ω_K be the maximal idempotent factor of μ . By a Theorem of Parthasarathy and Sazanov (see Theorem 7.2 of [4]), $\mu = \omega_K * \mu_1$ where μ_1 is an infinitely divisible measure without idempotent factors. Since T and S are automorphisms, $T(K)$ and $S(K)$ are the maximal idempotent factors of $T(\mu)$ and $S(\mu)$

respectively and $T(\mu_1)$ and $S(\mu_1)$ are infinitely divisible measures without idempotent factors. We also get that $\omega_K * \mu_1 = \omega_{T(K)S(K)} * T(\mu_1) * S(\mu_1)$. Since $T(\mu_1)$ and $S(\mu_1)$ are infinitely divisible without idempotent factors, by Theorem 4.2 of [5], the Fourier transform of $T(\mu_1)$ and $S(\mu_1)$ do not vanish. Let χ be any character of V . If $\hat{\omega}_K(\chi) = 0$, then $\hat{\omega}_{T(K)S(K)}(\chi) = 0$. If $\hat{\omega}_{T(K)S(K)}(\chi) = 0$, then since $\hat{\mu}_1(\chi) \neq 0$, we have $\hat{\omega}_K(\chi) = 0$. Thus, $\omega_K = \omega_{T(K)S(K)}$, that is, $K = T(K)S(K)$. Iteratively we prove that $\prod_{l(\alpha)=n} \alpha(K) = K$ for any $n \geq 1$. Since $\alpha(v) \rightarrow e$ as $l(\alpha) \rightarrow \infty$, $K \subset U$ for any compact open subgroup in V . Since V is totally disconnected, $K = (e)$. Thus, μ has no idempotent factors.

Theorem 4.1 *Let $T_1, T_2, \dots, T_m$ be contractions on a finite-dimensional vector space V over real or $\mathbb{Q}_p$ such that $T_i T_j = T_j T_i$ for all $1 \leq i, j \leq m$. Suppose $\mu = T_1(\mu) * T_2(\mu) * \dots * T_m(\mu) \in \mathcal{P}(V)$. Then μ is infinitely divisible without idempotent factors. In particular, autophage measures on V are infinitely divisible measures without idempotent factors.*

Proof We will treat the case when $m = 2$, the general case may be proved in a similar fashion. Let $k \geq 1$ be such that $\|T^k\| < 1$ and $\|S^k\| < 1$. Since T and S commute with each other, we get that $\mu = \prod_{i=0}^{2k} T^{2k-i} S^i(\mu)$. It is easy to see that $\|T^{2k-i} S^i\| < 1$ for any $0 \leq i \leq 2k$. By Proposition 4.1, we get that μ is an infinitely divisible measure without idempotent factors.

5 Absolute continuity

We now introduce a nondegeneracy condition which will be needed for absolute continuity. We say that a measure μ on a finite-dimensional vector space over real or $\mathbb{Q}_p$ is *full* if μ is not supported on a coset of a proper subspace. We now prove a result on the Fourier transform of full measures.

Lemma 5.1 *Let μ be a full measure on a finite-dimensional vector space V over real or $\mathbb{Q}_p$ and $\hat{\mu}$ be its Fourier transform. Suppose $\mu = T(\mu) * S(\mu)$ for some contractions T and S . Also, if T and S are strict-contractions or T and S commute with each other. Then $|\hat{\mu}(v)| \neq 1$ for all $v \in V \setminus (e)$.*

Proof Here we identify the dual of V with itself. Let $\lambda = \mu * \check{\mu}$. Then it is easy to see that

1. μ is full if and only if λ is not contained in a proper subspace,
2. for any $v \in V \setminus (e)$, $|\hat{\mu}(v)| = 1$ if and only if $\hat{\lambda}(v) = 1$.

In view of this it is enough to show that $\hat{\lambda}(v) \neq 1$ for any $v \in \setminus(e)$. Suppose $\hat{\lambda}(v) = 1$ for some $v \neq e$. Then the closed subgroup, say H , generated by the support of λ is a proper subgroup (in fact, λ is supported on the annihilator of the subgroup of all $v \in V$ such that $\hat{\lambda}(v) = 1 = |\hat{\lambda}(v)|$). By assumption, there exists contractions T and S such that $\lambda = T(\lambda) * S(\lambda)$. This implies that $T(H) \subset H$, since e is in the support of λ . Let H^0 be the maximal vector subspace contained in H . Then $T(H^0) \subset H^0$. Since T is an automorphism, $T(H^0) = H^0$. Similarly, we claim that $S(H^0) = H^0$. Let $\tilde{\lambda}$ denotes image of λ in V/H^0 and $\tilde{T}$ and $\tilde{S}$ be the factor automorphisms on V/H^0 corresponding to T and S . We now claim that $\tilde{\lambda} = \delta_e$. Suppose V is a real vector space. Then H/H^0 is discrete and hence countable. By Proposition 2.1, λ degenerates and hence $\lambda = \delta_e$. Suppose V is a p -adic vector space. Then $H/H^0 = K$ say, is a compact subgroup. Since $\tilde{\lambda}$ is autophage, we have $\tilde{T}(K)\tilde{S}(K) = K$. This implies that $\omega_K = T(\omega_K)S(\omega_K)$. By Proposition 4.1 or by Theorem 4.1, $K = (e)$, that is $\lambda = \delta_e$. This implies that μ is supported on a coset of a proper subspace H^0 .

Proposition 5.1 *Let $T_1, T_2, \dots, T_m$ be a finite-set of strict-contractions on a finite-dimensional vector space V over real or $\mathbb{Q}_p$. Suppose μ is a full probability measure on V such that $\mu = T_1(\mu) * T_2(\mu) * \dots * T_m(\mu)$. Then $\hat{\mu}$ is an integrable function.*

Proof We treat the case when $m = 2$, the general case may be proved in a similar fashion. Let T and S be two strict-contractions such that $\mu = T(\mu) * S(\mu)$. Let T^* and S^* be the adjoint of T and S respectively. Let $t = \|T^{*-1}\|^{-1}$ and $s = \|S^{*-1}\|^{-1}$. Let $0 < r < \infty$ be such that $t^r + s^r = 1$.

Let $\hat{\mu}$ be the Fourier transform of μ . By Theorem 4.1 and Theorem 4.2 of [5], $\hat{\mu}$ does not vanish. Thus, the function g defined by

$$g(v) = -\|v\|^{-r} \log(|\hat{\mu}(v)|)$$

for any $v \in V \setminus (e)$ is a continuous function. Since μ is full, by Lemma 5.1, $|\hat{\mu}(v)| \neq 1$ for any $v \in V \setminus (e)$. Thus, g does not vanish.

Now for any $v \in V \setminus (e)$,

$$\begin{aligned} g(v) &= -\|v\|^{-r} [\log |\hat{\mu}(T^*(v))| + \log |\hat{\mu}(S^*(v))|] \\ &= \left(\frac{\|T^*(v)\|}{\|v\|}\right)^r g(T^*(v)) + \left(\frac{\|S^*(v)\|}{\|v\|}\right)^r g(S^*(v)) \\ &\geq t^r g(T^*(v)) + s^r g(S^*(v)) \end{aligned}$$

since for any automorphism α of V and $v \in V \setminus (e)$, $\|v\| \leq \|\alpha^{-1}\| \|\alpha(v)\|$. Thus, for any $v \neq 0$ in V , we have $g(v) \geq \min\{g(T^*(v)), g(S^*(v))\}$. For each $v \in V \setminus (e)$, we now define for $n \geq 0$, $\alpha_n \in sg(T^*, S^*)$ inductively as follows: $\alpha_0 = I$ and for $n \geq 1$, α_n to be the either $T^*\alpha_{n-1}$ or $S^*\alpha_{n-1}$ if $g(T^*\alpha_{n-1}(v)) \leq g(S^*\alpha_{n-1}(v))$ or

$g(S^*\alpha_{n-1}(v)) \leq g(T^*\alpha_{n-1}(v))$ respectively. This implies that

$$\begin{aligned} g(\alpha_{n-1}(v)) &\geq t^r g(T^*\alpha_{n-1}(v)) + s^r g(S^*\alpha_{n-1}(v)) \\ &\geq \min\{g(T^*\alpha_{n-1}(v)), g(S^*\alpha_{n-1}(v))\} \\ &= g(\alpha_n(v)) \end{aligned}$$

for all $n \geq 1$. Thus, for each $v \in V \setminus (e)$, there exists a sequence (α_n) (possibly, depending on v) in $sg(T^*, S^*)$ such that $l(\alpha_n) = n$, $\alpha_n = T^*\alpha_{n-1}$ or $S^*\alpha_{n-1}$ and $g(v) \geq g(\alpha_n(v))$.

Let $c = \min\{g(v) \mid \min\{t, s\} \leq \|v\| \leq 1\}$. We now claim that $0 < c \leq g(v)$ for any $\|v\| > 1$. Since g does not vanish, $c > 0$. Let v be such that $\|v\| > 1$. Let $\alpha_n \in sg(T^*, S^*)$ be such that $l(\alpha_n) = n$, $\alpha_n = T^*\alpha_{n-1}$ or $S^*\alpha_{n-1}$ and $g(\alpha_n(v)) \leq g(v)$. Then we get that $\alpha_n(v) \rightarrow 0$ as $n \rightarrow \infty$ and $g(v) \geq g(\alpha_n(v))$ for all n . Let $k \in \mathbb{N}$ be such that $\|\alpha_k(v)\| \leq 1$ but $\|\alpha_{k-1}(v)\| > 1$. Since the roles of T and S are similar, we may assume that $\alpha_k = T^*\alpha_{k-1}$. So,

$$1 < \|\alpha_{k-1}(v)\| \leq \|T^{*-1}\| \|\alpha_k(v)\|$$

and hence $t = \|T^{*-1}\|^{-1} \leq \|\alpha_k(v)\| \leq 1$. This shows that

$$g(v) \geq c$$

for all v with $\|v\| > 1$. It follows from the definition of g that

$$|\hat{\mu}(v)| \leq \exp(-c\|v\|^r)$$

for all v with $\|v\| > 1$ and for some constant $c > 0$. Thus, $\hat{\mu}$ is an integrable function.

Theorem 5.1 *Let $T_1, T_2, \dots, T_m$ be contractions on a finite-dimensional vector space V over real or $\mathbb{Q}_p$ such that $T_i T_j = T_j T_i$ for all $1 \leq i, j \leq m$. Suppose $\mu = T_1(\mu) * T_2(\mu) * \dots * T_m(\mu) \in \mathcal{P}(V)$ is full. Then μ is absolutely continuous with a bounded continuous density. In particular, a full autophage measure on V is absolutely continuous with a bounded continuous density.*

Remark 5.1 Any question regarding absolute continuity should be addressed only in the class of full measures because any non-full measure is obviously not absolutely continuous. In view of this the assumption fullness is necessary in Proposition 5.1 and Theorem 5.1.

Proof Arguing as in Theorem 4.1, we may assume that each T_i is strict-contraction. Now by Proposition 5.1, we get that $\hat{\mu}$, the Fourier transform of μ is integrable. This shows by inverse Fourier transform that the measure μ is absolutely continuous with a bounded continuous density.

We now prove the absolute continuity of certain semistable measures.

Corollary 5.1 *Let μ be a full measure on a finite-dimensional vector space over real or $\mathbb{Q}_p$. Suppose μ is (T, n) -semistable. Then μ is absolutely continuous with a bounded continuous density.*

Proof Suppose μ is (T, n) -semistable, that is, $T(\mu^n) = \mu$. Suppose V is a real vector space, then since μ is full by [2], we get that T is a contraction. By Theorem 5.1, μ is absolutely continuous with a bounded continuous density. Suppose V is a vector space over $\mathbb{Q}_p$. Then by [8], there exist a $x \in V$ and a continuous convolution semigroup (μ_t) in $\mathcal{P}(V)$ such that $T(\mu_t) = \mu_{nt}$ for all $t > 0$ and $\mu_1 = x\mu$. It is enough to prove μ_1 is absolutely continuous. By [1], there exist a compact subgroup K of V and a convolution semigroup $\tilde{\mu}_t$ in $\mathcal{P}(C(T))$ where $C(T) = \{x \in V \mid T^n(x) \rightarrow e\}$ such that $\mu_t = \omega_k * \tilde{\mu}_t$, $T(\tilde{\mu}_t) = \tilde{\mu}_{nt}$ and $C(T) \cap K = (e)$. Since μ is full, V is spanned by $C(T)$ and K . Let W be the vector space spanned by K . Then K is open in W and $C(T) \cap W = (e)$. Thus, V is the direct product of $C(T)$ and W . Since $\mu_1 = \omega_K * \tilde{\mu}_1$, it is enough to show that $\tilde{\mu}_1$ is absolutely continuous. Since μ is full, μ_1 is full and hence $\tilde{\mu}_1$ is full in $C(T)$. By Theorem 5.1, $\tilde{\mu}_1$ is absolutely continuous with a bounded continuous density.

Acknowledgement I wish to thank ICTP, Trieste, Italy for providing financial support and all facilities during my stay at ICTP. I also wish to thank NBHM, India for providing travel grant to visit ICTP.

References

- [1] S. G. Dani and R. Shah, Contraction subgroups and semistable measures on p -adic Lie groups, Math. Proc. Cambridge Philos. Soc. 110 (1991), 299-306.
- [2] W. Hazod, Remarks on (semi-)stable probabilities, Probability measures on groups, VII (Oberwolfach, 1983), Lecture Notes in Math., 1064, Springer, Berlin, 1984, 182-203.
- [3] A. Luczak, Operator semistable probability measures on R^N , Colloq. Math. 45 (1981), 287-300.
- [4] K. R. Parthasarathy, Probability measures on metric spaces, Probability and Mathematical Statistics, No.3, Academic Press, Inc., New York-London 1967.
- [5] K. R. Parthasarathy, R. Ranga Rao and S. R. S. Varadhan, Probability distributions on locally compact abelian groups, Illinois J. math. 7 (1963), 337-369.
- [6] C. R. E. Raja and Riddhi Shah, Factorization of T -decomposable measures on groups, Monatsh. Math. 133 (2001), 223-239.

- [7] I. Z. Rusza and G. J. Szekely, Algebraic Probability Theory, Wiley Series in Probability in Mathematics and Statistics, John Wiley and Sons, Chichester-New York-Singapore, 1988.
- [8] R. Shah, Semistable measures and limit theorems on real and p -adic groups, Monatsh. Math. 115 (1993), 191-213.
- [9] E. Siebert, Strongly operator-decomposable probability measures on separable Banach spaces, Math. Nachr. 154 (1991), 315-326.
- [10] G. J. Szekely, Autophage and allelophage probability distributions, Probability theory and mathematical statistics with applications (Visegrád, 1985), 205-211.

Present address: Mathematics Section, ICTP, Strada Costiera 11, 34014 Trieste. Italy.

e-amil: creraja@ictp.trieste.it

Permanent address: Stat-Math Unit, Indian Statistical Institute, 8th Mile Mysore Road, R. V. College Post, Bangalore - 560 059. India.

e-amil: creraja@isibang.ac.in